

# **Certificate of Analysis**

www.tocris.com

Product Name: 8-Chloroadenosine Catalog No.: 4436 Batch No.: 2

CAS Number: 34408-14-5

IUPAC Name: 6-Amino-8-chloropurine riboside

# 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula:  $C_{10}H_{12}CIN_5O_4.H_2O$ 

Batch Molecular Weight: 319.71

Physical Appearance: White solid

**Solubility:** water to 20 mM with gentle warming

DMSO to 100 mM

Storage: Store at -20°C

Batch Molecular Structure:

# 2. ANALYTICAL DATA

HPLC: Shows 98.6% purity

1H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 37.57 4.41 21.91 Found 37.74 4.23 21.76



# **Product Information**

Print Date: May 10th 2013

www.tocris.com

Product Name: 8-Chloroadenosine Catalog No.: 4436 Batch No.: 2

CAS Number: 34408-14-5

IUPAC Name: 6-Amino-8-chloropurine riboside

#### **Description:**

Nucleoside analog; metabolized in vivo to 8-Chloro-ATP. Incorporates into RNA during transcription and inhibits RNA synthesis. Exhibits cytotoxicity in MM.1S, RPMI-8226 and U266 cancer cell lines; induces  $G_2/M$  cell cycle arrest and mitotic catastrophe in A549 and H1299 cells.

# **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>10</sub>H<sub>12</sub>ClN<sub>5</sub>O<sub>4</sub>.H<sub>2</sub>O

Batch Molecular Weight: 319.71 Physical Appearance: White solid

Minimum Purity: >98%

#### **Batch Molecular Structure:**

Storage: Store at -20°C

## Solubility & Usage Info:

water to 20 mM with gentle warming DMSO to 100 mM

## Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

## References:

**Zhang** *et al* (2004) Exposure of human lung cancer cells to 8-chloro-adenosine induces G2/M arrest and mitotic catastrophe. Neoplasia *6* 802. PMID: 15720807.

**Gu** et al (2006) 8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells Biochem.Pharmacol. **72** 541. PMID: 16844099.

**Cervantes-Gomez** *et al* (2011) ATP analog enhances the actions of a heat shock protein 90 inhibitor in multiple myeloma cells. J.Pharmacol.Exp.Ther. **339** 545. PMID: 21821695.

