

Certificate of Analysis

www.tocris.com

Product Name: TAME hydrochloride

CAS Number: **IUPAC Name:**

Catalog No.: 4506 EC Number: 217-235-1 Batch No.: 1


1784-03-8

N²-[(4-Methylphenyl)sulfonyl]-L-arginine methyl ester hydrochloride

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: Batch Molecular Weight: Physical Appearance: Solubility:

Storage: **Batch Molecular Structure:** C14H22N4O4S.HCI 378.87 White solid water to 100 mM DMSO to 100 mM Store at -20°C



2. ANALYTICAL DATA

Melting Point: HPLC: ¹H NMR: Mass Spectrum: **Optical Rotation: Microanalysis:**

At 149°C Shows 100% purity Consistent with structure Consistent with structure $[\alpha]_D = -14.1$ (Concentration = 4, Solvent = Water) Carbon Hydrogen Nitrogen Theoretical 44.38 6.12 14.79 Found 44.4 6.13 14.73

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use





Product Information

Print Date: Apr 28th 2015

www.tocris.com

Product Name: TAME hydrochloride

CAS Number: **IUPAC Name:**

1784-03-8

Catalog No.: 4506

Batch No.: 1

EC Number: 217-235-1

 N^{2} -[(4-Methylphenyl)sulfonyl]-L-arginine methyl ester hydrochloride

Description:


Competitive inhibitor of the ubiquitin ligase anaphase-promoting complex/cyclosome (APC/C); promotes Cdc20 autoubiquitination. Inhibits cyclin proteolysis in mitotic Xenopus oocyte extracts (IC₅₀ = 12 μ M) and blocks cyclin degradation in interphase extract.

Physical and Chemical Properties:

Batch Molecular Formula: C14H22N4O4S.HCI Batch Molecular Weight: 378.87 Physical Appearance: White solid

Minimum Purity: >99%

Batch Molecular Structure:



Storage: Store at -20°C

Solubility & Usage Info: water to 100 mM DMSO to 100 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Zeng et al (2010) Pharmacologic inhibition of the anaphase-promoting complex induces a spindle checkpoint-dependent mitotic arrest in the absence of spindle damage. Cancer Cell. 18 382. PMID: 20951947.

Zeng & King (2012) An APC/C inhibitor stabilizes cyclin B1 by prematurely terminating ubiquitination. Nat.Chem.Biol. 8 383. PMID: 22366722.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

