

Certificate of Analysis

www.tocris.com

Product Name: VUF 11207 fumarate Catalog No.: 4780 Batch No.: 1

IUPAC Name: N-[(2E)-3-(2-Fluorophenyl)-2-methyl-2-propen-1-yl]-3,4,5-trimethoxy-<math>N-[2-(1-methyl-2-pyrrolidinyl)]

benzamide fumarate

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{27}H_{35}FN_2O_4.C_4H_4O_4.H_2O_4$

Batch Molecular Weight: 604.67 **Physical Appearance:** White solid

Solubility: water to 100 mM

DMSO to 100 mM

Storage: Store at -20°C

Batch Molecular Structure:

2. ANALYTICAL DATA

TLC: $R_f = 0.4$ (Chloroform:Methanol:Ammonia soln. [95:5:0.1])

HPLC: Shows >99.3% purity
¹H NMR: Consistent with structure
Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 61.58 6.83 4.63 Found 61.57 6.48 4.63



Product Information

Print Date: Mar 8th 2014

www.tocris.com

Product Name: VUF 11207 fumarate Catalog No.: 4780 Batch No.: 1

IUPAC Name: N-[(2E)-3-(2-Fluorophenyl)-2-methyl-2-propen-1-yl]-3,4,5-trimethoxy-<math>N-[2-(1-methyl-2-pyrrolidinyl)]

benzamide fumarate

Description:

Potent CXCR7 chemokine receptor agonist (EC $_{50}$ = 1.6 nM). Induces recruitment of β -arrestin to CXCR7 in HEK293-CXCR7 cells.


Physical and Chemical Properties:

Batch Molecular Formula: C₂₇H₃₅FN₂O₄.C₄H₄O₄.H₂O

Batch Molecular Weight: 604.67 Physical Appearance: White solid

Batch Molecular Structure:

Minimum Purity: >98%



Storage: Store at -20°C

Solubility & Usage Info:

water to 100 mM DMSO to 100 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Wijtmans et al (2012) Synthesis, modeling and functional activity of substituted styrene-amides as small-molecule CXCR7 agonists. Eur.J.Med.Chem. 51 184. PMID: 22424612.

