

Certificate of Analysis

www.tocris.com

Product Name: PA 452 Catalog No.: 5086 Batch No.: 1

CAS Number: 457657-34-0

IUPAC Name: 2-[[3-(Hexyloxy)-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl]methylamino]-5-pyrimidinecarboxylic

acid

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{26}H_{37}N_3O_3$. ¹/₄H₂O

Batch Molecular Weight: 444.09

Physical Appearance: Off-white solid
Solubility: DMSO to 100 mM

ethanol to 10 mM

Storage: Store at +4°C

Batch Molecular Structure:

2. ANALYTICAL DATA

TLC: $R_f = 0.32$ (Ethyl acetate:Petroleum ether [1:1])

HPLC: Shows >99.8% purity
 ¹H NMR: Consistent with structure
 Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 70.32 8.51 9.46 Found 70.38 8.3 9.58



Product Information

Print Date: May 29th 2014

www.tocris.com

Product Name: PA 452 Catalog No.: 5086 Batch No.: 1

CAS Number: 457657-34-0

IUPAC Name: 2-[[3-(Hexyloxy)-5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl]methylamino]-5-pyrimidinecarboxylic

acid

Description:

RXR antagonist (pA $_2$ = 7.11). Triggers dissociation of RXR tetramers. Attenuates cell proliferation and induces apoptosis in MCF-7 breast cancer cells.

Physical and Chemical Properties:

Batch Molecular Formula: C₂₆H₃₇N₃O₃. ¼H₂O

Batch Molecular Weight: 444.09 Physical Appearance: Off-white solid

Minimum Purity: >98%

Batch Molecular Structure:

Storage: Store at +4°C

Solubility & Usage Info:

DMSO to 100 mM ethanol to 10 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Takahashi et al (2002) Novel retinoid X receptor antagonists: specific inhibition of retinoid synergism in RXR-RAR heterodimer actions. J.Med.Chem. **45** 3327. PMID: 12139443.

Yasmin et al (2010) Inhibition of mammary carcinoma cell growth by RXR is mediated by the receptor's oligomeric switch. J.Mol.Biol. 397 1121. PMID: 20188110.

Nakayama *et al* (2011) Discovery of a potent retinoid X receptor antagonist structurally closely related to RXR agonist NEt-3IBIB. J.Med.Chem.Lett **2** 896.

