

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD68/SR-D1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD68/SR-D1 Asn22-Ile320 Accession # AAB25811
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human CD68/SR-D1

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CD68, also called Scavenger Receptor D1 (SR-D1) or macrophage scavenger receptor 1, is a 110 kDa type I transmembrane glycoprotein that belongs to the LAMP family of molecules (1). It contains a 300 amino acid (aa) extracellular region that is rich in threonine and serine, a likely attachment site for multiple carbohydrates. Human CD68 shares 74% aa sequence identity to mouse CD68 in the extracellular region. CD68 is found on monocytes and macrophages and serves as a scavenger receptor for oxidized LDL.

References:

1. Holness, C.L. and D.L. Simmons (1993) *Blood* **81**:1607.