

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse S100A9 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 2% cross-reactivity with recombinant mouse S100A10 and recombinant human S100B is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse S100A9 Ala2-Lys113 Accession # P31725
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse S100A9
Immunocytochemistry	5-15 µg/mL	See Below

DATA

Immunocytochemistry



S100A9 in XB2 Mouse Cell Line. S100A9 was detected in immersion fixed XB2 mouse teratoma keratinocyte cell line using 10 µg/mL Goat Anti-Mouse S100A9 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF2065) for 3 hours at room temperature. Cells were stained with the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (red; Catalog # NL001) and counterstained with DAPI (blue). View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

S100A9, also known as calgranulin B, migration inhibitory factor-related protein 14 (MRP14), belongs to the S100 family of calcium binding proteins of the EF-hand type. It is highly expressed in myeloid cells and other cell types including keratinocytes, epithelial cells and endothelial cells. S100A9 is a cytosolic protein that can also be released into the extracellular milieu. It is associated with various inflammatory diseases.