



## ***Anti-rat EG-VEGF/PK1 Antibody***

### **ORDERING INFORMATION**

**Catalog Number:** AF2100

**Lot Number:** KUC01

**Size:** 100 µg

**Formulation:** 0.2 µm filtered solution in PBS  
with 5% trehalose

**Storage:** -20° C

**Reconstitution:** sterile PBS

**Specificity:** rat, mouse and human EG-VEGF

**Immunogen:** NS0-derived rrEG-VEGF

**Ig Type:** sheep IgG

**Applications:** Direct ELISA  
Western blot

### ***Preparation***

Produced in sheep immunized with purified, NS0-derived, recombinant rat Endocrine Gland-derived Vascular Endothelial Growth Factor (rrEG-VEGF). EG-VEGF specific IgG was purified by rat EG-VEGF affinity chromatography.

### ***Formulation***

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) with 5% trehalose.

### ***Endotoxin Level***

< 0.1 EU per 1 µg of the antibody as determined by the LAL method.

### ***Reconstitution***

Reconstitute with sterile PBS. If 0.5 mL of PBS is used, the antibody concentration will be 0.2 mg/mL.

### ***Storage***

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

### ***Specificity***

This antibody has been selected for its ability to recognize rat, mouse and human EG-VEGF in direct ELISAs and western blots.

### ***Applications***

**Direct ELISA** - This antibody can be used at 0.5 - 1.0 µg/mL with the appropriate secondary reagents to detect rat, mouse and human EG-VEGF. The detection limit for rrEG-VEGF, rmEG-VEGF and rhEG-VEGF is approximately 0.7 ng/well.

**Western blot** - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect rat, mouse and human EG-VEGF. The detection limit for rrEG-VEGF, rmEG-VEGF and rhEG-VEGF is approximately 1 ng/lane under non-reducing and reducing conditions.

**Optimal dilutions should be determined by each laboratory for each application.**