

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects endogenous human and mouse BCAP in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human BCAP Met462-Leu650 Accession # Q6ZUJ8
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

BCAP (B-cell adaptor for phosphoinositide 3-kinase (PI3K)) participates in linking the B cell antigen receptor (BCR) with the PI3K pathway. Tyrosine phosphorylation of BCAP by BCR-associated protein tyrosine kinases, such as Syk and Btk, generates binding sites for the p85 subunit of PI3K, resulting in activation of the PI3K pathway. Human and mouse BCAP contain 3 YXXM SH2 binding motifs for interaction with PI3K.