



## ***Anti-rat CINC-2 $\alpha$ $\beta$ Antibody***

### **ORDERING INFORMATION**

**Catalog Number:** AF517

**Lot Number:** BFA01

**Size:** 100  $\mu$ g

**Formulation:** 0.2  $\mu$ m filtered solution in PBS

**Storage:** -20° C

**Reconstitution:** sterile PBS

**Specificity:** rat CINC-2 $\alpha$  and rat CINC-2 $\beta$

**Immunogen:** *E. coli*-derived rrCINC-2 $\beta$

**Ig Type:** rat CINC-2 $\alpha$  and CINC-2 $\beta$  specific  
goat IgG

**Applications:** Neutralization of bioactivity  
Western blot  
ELISA

### ***Preparation***

Produced in goats immunized with purified, *E. coli*-derived, recombinant rat cytokine-induced neutrophil chemoattractant 2 beta (rrCINC-2 $\beta$ ). CINC-2 $\alpha$ / $\beta$  specific IgG was purified by rat CINC-2 $\beta$  affinity chromatography.

### ***Formulation***

Lyophilized from a 0.2  $\mu$ m filtered solution in phosphate-buffered saline (PBS).

### ***Endotoxin Level***

< 0.1 EU per 1  $\mu$ g of the antibody as determined by the LAL method.

### ***Reconstitution***

Reconstitute with sterile PBS. If 1 mL of PBS is used, the antibody concentration will be 0.1 mg/mL.

### ***Storage***

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

### ***Specificity***

This antibody has been selected for its ability to neutralize rrCINC-2 $\beta$  bioactivity. Based on direct ELISA results, this antibody shows almost 100% cross-reactivity with rrCINC-2 $\alpha$ , approximately 50% cross-reactivity with rmMIP-2, 35% cross-reactivity with rmKC, 15% cross-reactivity with rrCINC-1 and 10% cross-reactivity with rhIL-8, rhGRO $\alpha$ , rhGRO $\beta$  and rhGRO $\gamma$ . Additionally, in direct ELISA, this antibody shows no cross-reactivity with other chemokines tested.<sup>1</sup>

### ***Neutralization of Rat CINC-2 $\beta$ Bioactivity***

The exact concentration of antibody required to neutralize rrCINC-2 $\beta$  activity is dependent on the cytokine concentration, cell type, growth conditions and the type of activity studied. To provide a guideline, R&D Systems has determined the neutralization dose for this antibody under a specific set of conditions. The **Neutralization Dose<sub>50</sub> (ND<sub>50</sub>)** for this antibody is defined as that concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when that cytokine is present at a concentration just high enough to elicit a maximum response.

The ND<sub>50</sub> for this lot of anti-rat CINC-2 $\beta$  antibody was determined to be approximately 0.4 - 2  $\mu$ g/mL in the presence of 0.05 ng/mL of rrCINC-2 $\beta$ , using BaF/3 cells transfected with rhCXCR-2. The specific conditions are described in the figure legends.

### ***Additional Applications***

**Western blot** - This antibody can be used at 0.1 - 0.2  $\mu$ g/mL with the appropriate secondary reagents to detect rat CINC-2 $\beta$ . The detection limit for rrCINC-2 $\beta$  is approximately 5 ng/lane and 25 ng/lane under non-reducing and reducing conditions, respectively.

**Direct ELISA** - This antibody can be used at 0.5 - 1.0  $\mu$ g/mL with the appropriate secondary reagents to detect rat CINC-2 $\beta$ . The detection limit for rrCINC-2 $\beta$  is approximately 0.5 ng/well.


**Optimal dilutions should be determined by each laboratory for each application.**

FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

**R&D Systems, Inc.**  
**1-800-343-7475**


**Figure 1**

**Chemotactic Effect of rrCINC-2 $\beta$**



**Figure 2**

**Neutralization of rrCINC-2 $\beta$**



**Figure 1**

Recombinant rat CINC-2 $\beta$  chemoattracts BaF/3 cells transfected with rhCXCR-2. The ED<sub>50</sub> for this effect is typically 5 - 25 ng/mL.

**Figure 2**

To measure the ability of the antibody to neutralize the chemoattractant activity of rrCINC-2 $\beta$  for hCXCR-2 transfected BaF/3 cells, rrCINC-2 $\beta$  was incubated with various concentrations of the antibody for 15 minutes at room temperature in a 96 well microplate. Following this preincubation period, 75  $\mu$ L of the cytokine-antibody solution (containing rrCINC-2 $\beta$  at a final concentration of 0.05  $\mu$ g/mL and antibody at the concentrations indicated) was transferred to the lower compartment of a 96 well chemotaxis chamber (NeuroProbe, Cabin John, MD). The chemotaxis chamber was then assembled using a PVP-free polycarbonate filter (5 micron pore size) and  $0.2 \times 10^6$  cells/well was added to the top chamber. After incubation for 3 hours at 37 $^\circ$  C in a 5% CO<sub>2</sub> humidified incubator, the chamber was carefully disassembled. The cells that migrate through to the lower chamber were transferred to a 96 well plate. Chemotaxis was measured by Resazurin (R&D Systems, Catalog # AR002) staining of cells that have migrated through the filter. As shown in figure 2, the ND<sub>50</sub> for this lot of antibody is approximately 0.4 - 2  $\mu$ g/mL.