

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Mouse |
| Specificity | Detects mouse CD117/c-kit in Western blots. In Western blots, approximately 10% cross-reactivity with recombinant human CD117 is observed. |
| Source | Polyclonal Goat IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant mouse CD117/c-kit Gln25-Thr519 (Ala207Glu) Accession # P05532 |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------------|----------------------------------|---|
| Western Blot | 0.1 µg/mL | Recombinant Mouse CD117/c-kit |
| Flow Cytometry | 0.25 µg/10 ⁶ cells | Lineage depleted mouse bone marrow |
| Immunohistochemistry | 5-15 µg/mL | Immersion fixed frozen sections of mouse embryo (E15) |

PREPARATION AND STORAGE

| | |
|--------------------------------|---|
| Reconstitution | Reconstitute at 0.2 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | <p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

Stem cell factor receptor (CD117, the gene product of the c-kit protooncogene) and its ligand, stem cell factor (also named c-kit ligand, mast cell growth factor), play essential roles in gametogenesis, melanogenesis and hematopoiesis. It is a transmembrane tyrosine kinase that is expressed on endothelial cells, mast cells, megakaryocytes, stem cells and multiple embryonic cells, such as melanoblasts and primordial germ cells.