



## ***Biotinylated Anti-human CRISP-3 Antibody***

### **ORDERING INFORMATION**

**Catalog Number:** BAF2397

**Lot Number:** UON01

**Size:** 50 µg

**Formulation:** 0.2 µm filtered solution in PBS  
with BSA

**Storage:** -20° C

**Reconstitution:** sterile 0.1% BSA in TBS

**Specificity:** human CRISP-3

**Immunogen:** NS0-derived rhCRISP-3

**Ig Type:** goat IgG

**Application:** Western blot

### ***Preparation***

Produced in goats immunized with purified, NS0-derived, recombinant human Cysteine-Rich Secretory Protein-3 (rhCRISP-3). Human CRISP-3 specific IgG was purified by human CRISP-3 affinity chromatography and then biotinylated. CRISP-3 belongs to the large CRISP family of secreted proteins expressed in mammals, reptiles, insects and plants. All family members share a common cystein-rich domain at the carboxy-terminus. Human CRISP-3 is found in salivary glands, prostate, pancreas and in secretory granules of neutrophilic granulocytes. In plasma, human CRISP-3 is complexed with alpha (1) B-glycoprotein, a member of the Ig superfamily. Human CRISP-3 shares 71% and 48% amino acid sequence identity with human CRISP-2 and mouse CRISP-3, respectively.

### ***Formulation***

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) containing 50 µg of bovine serum albumin (BSA) per 1 µg of antibody.

### ***Reconstitution***

Reconstitute with sterile Tris-buffered saline pH 7.3 (20 mM Trizma base, 150 mM NaCl) containing 0.1% BSA. If 1 mL of buffer is used, the antibody concentration will be 50 µg/mL.

### ***Storage***

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

### ***Specificity***

This antibody has been selected for use as a detection antibody in human CRISP-3 western blots.

### ***Application***

**Western Blot** - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect human CRISP-3. The detection limit for rhCRISP-3 is approximately 5 ng/lane under non-reducing and reducing conditions. In this format, this antibody shows approximately 15% cross-reactivity with rhCRISP-2.

**Optimal dilutions should be determined by each laboratory for each application.**