

## DESCRIPTION


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects Gr-1/Ly-6G. Weak cross-reactivity with Ly-6C is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # RB6-8C5
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Conjugate</b>	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 µL/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> ● 12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

The myeloid differentiation antigen Gr-1 is a member of the Ly-6 family, Ly-6G (1). The RB6-8C5 antibody also reacts weakly with Ly-6C-transfected EL4 cells (2). In the periphery, this antibody specifically recognized granulocytes (3, 4).

### References:

1. Spangrude, G.J. *et al.* (1988) *Science* **241**:58.
2. Fleming, T.J. *et al.* (1993) *J. Immunol.* **151**:2399.
3. Lewinsohn, D.M. *et al.* (1987) *J. Immunol.* **147**:22.
4. Lagasse, E. and I.L. Weissman (1996) *J. Immunol. Methods* **197**:139.