

# Mouse Integrin αM/CD11b PerCP-conjugated Antibody

Monoclonal Rat IgG<sub>2B</sub> Clone # M1/70

Catalog Number: FAB1124C

-	-		-		-	_
	_				_	_
1	()	()	Iŀ	ΞS	П	۶.


DESCRIPTION					
Species Reactivity	Mouse				
Specificity	Detects mouse Integrin αM/CD11b. Cross-reaction with human Integrin αM has been reported (1, 2).				
Source	Monoclonal Rat IgG <sub>2B</sub> Clone # M1/70				
Purification	Protein A or G purified from hybridoma culture supernatant				
Immunogen	Con A-activated C57BL/10 splenocytes				
Conjugate	PerCP (Peridinin-chlorophyll Protein Complex) Excitation Wavelength: 482 and 564 nm Emission Wavelength: 675 nm				
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.				
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.				

#### **APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 <sup>6</sup> cells	See Below

#### DATA



Detection of Integrin aM/CD11b in Mouse Splenocytes by Flow Cytometry. Mouse splenocytes were stained with Rat Anti-Mouse Integrin aM/CD11b PerCPconjugated Monoclonal Antibody (Catalog # FAB1124C) and Rat Anti-Mouse F4/80/EMR1 PE-conjugated Monoclonal Antibody (Catalog # FAB5580P). Quadrant markers were set based on control antibody staining (Catalog # IC013C). View our protocol for Staining Membrane-associated Proteins.

#### PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below

Stability & Storage

Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

The Integrin family proteins are heterodimeric transmembrane receptors composed of an α and a β subunit. The Integrin αM subunit, also known as MAC-1α subunit or CD11b, combines with the Integrin β2 subunit (CD18) to form the non-covalent heterodimer Integrin αM/β2, also known as MAC-1 and Complement Receptor type 3 (CR3). Integrin αM/β2 is expressed on granulocytes, macrophages, dendritic cells and natural killer cells. Upon activation, αM/β2 can bind several ligands (including ICAM-1, Fibrinogen, and the C3 complement fragment, C3bi) to mediate phagocyte adhesion, migration and ingestion of complement-opsonized particles

### References:

- 1. Beller, D.J. et al. (1982) J. Exp. Med. 156:1000.
- Ault, K.A. and T.A. Springer (1981) J. Immunol. 126:359.

