

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Mouse |
| Specificity | Detects mouse Rae-1 ϵ in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse Rae-1 α , β , γ , or δ is observed. By flow cytometry, no cross-reactivity with mouse Rae-1 α or mouse Rae-1 γ . |
| Source | Monoclonal Rat IgG _{2A} Clone # 205001 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | BaF3 mouse pro-B cell line transfected with mouse Rae-1 ϵ |
| Conjugate | Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|--------------------------------------|--|
| Flow Cytometry | 0.25-1 μ g/10 ⁶ cells | BaF3 mouse pro-B cell line transfected with mouse Rae-1 ϵ |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

Rae-1 ϵ is a member of a family of cell-surface proteins that function as ligands for mouse NKG2D. Other family members are designated Rae-1 α , β , γ , and δ . Amino acid sequence identity within this family ranges from 88-95%. The Rae-1 proteins are distantly related to MHC class I proteins, but they possess only the α 1 and α 2 Ig-like domains, and they have no capacity to bind peptide or interact with β 2-microglobulin. The genes encoding these proteins are not found within the Major Histocompatibility Complex on mouse chromosome 17, but rather map to mouse chromosome 10. The Rae-1 proteins are anchored to the membrane via a GPI-linkage. The name of this family derives from the original identification of these proteins as the product of retinoic acid early inducible transcripts. Rae-1 expression is developmentally controlled. Transcripts were observed in the brain/head region of day 10-14 embryos but disappeared by day 18. Rae-1 transcripts were detected in several transformed cell lines but are absent from most normal adult tissues. All Rae-1 family members bind to mouse NKG2D, an activating receptor expressed on NK cells and some T cell subsets, resulting in the activation of cytolytic activity and/or cytokine production by these effector cells. Ectopic expression of Rae-1 on mouse tumor cell lines resulted in the *in vivo* rejection of the tumors (1-7).

References:

1. Zou, Z. *et al.* (1996) *J. Biochem (Tokyo)* **119**:319.
2. Diefenbach, A. *et al.* (2000) *Nature Immunol.* **1**:119.
3. Cerwenka, A. *et al.* (2000) *Immunity* **12**:721.
4. Cerwenka, A. *et al.* (2001) *Proc. Natl. Acad. Sci. USA* **98**:11521.
5. Diefenbach, A. *et al.* (2001) *Nature* **413**:165.
6. Champsaur, M. *et al.* (2010) *J. Immunol.* **185**:157.
7. Markiewicz M. *et al.* (2012) *Immunity* **36**:132.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.