

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Pref-1/DLK1/FA1 in direct ELISAs and Western blots. Shows approximately 50% cross-reactivity with recombinant mouse Pref-1/DLK1/FA1.
Source	Monoclonal Mouse IgG _{2B} Clone # 211309
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Pref-1/DLK1/FA1 long isoform Ala24-Pro297 (with Arg248Pro and Lys295Ser substitutions) Accession # P80370
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Pref-1 (Preadipocyte Factor 1), also known as DLK1 and FA1, is a 58-65 kDa member of the Notch/Serrata/Delta family of proteins. It is expressed in prechondrocytes and preadipocytes and appears to block progenitor cell differentiation into mature cell lineages. Mature human Pref-1 is a 360 amino acid (aa) type I transmembrane N- and O-linked glycoprotein. It contains a 280 aa extracellular region (aa 24-303), a 24 aa transmembrane segment (aa 304-327), and a 56 aa cytoplasmic domain (aa 328-383). The extracellular region contains six EGF-like domains, and undergoes proteolytic cleavage to generate a bioactive 50 kDa fragment, plus three 25-31 kDa fragments that show no activity. There are multiple potential splice variants. One shows a deletion of aa 229-301, a second possesses a six aa substitution for aa 1-52, a third shows a deletion of aa 210-277, while a fourth contains a six aa substitution for aa 207-383. Over aa 24-297, human Pref-1 shares 82% aa identity with mouse Pref-1.