

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse H60 in ELISAs and Western blots. In flow cytometry, stains mouse H60 transfectants but not irrelevant transfectants.
Source	Monoclonal Rat IgG _{2A} Clone # 205326
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	BaF3 mouse pro-B cell line transfected with mouse H60 and mouse myeloma cell line NS0-derived recombinant mouse H60 Asp30-Gln212 Accession # Q3TDZ7
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	5 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

H60 was originally described as an immunodominant histocompatibility antigen that is expressed in BALB mice but not in B6 mice. More recently it was reported to function as a ligand for mouse NKG2D, an activating receptor found on NK cells, on some T cell subsets, and on stimulated macrophages. H60 shares approximately 25 percent amino acid identity with the Rae-1 family, a small group of proteins that also function as ligands for mouse NKG2D. H60 and the Rae-1 proteins are distantly related to MHC class I proteins, but they possess only the α1 and α2 Ig-like domains, and have no capacity to bind peptide or interact with β2-microglobulin. The genes encoding these proteins are not found within the Major Histocompatibility Complex on mouse chromosome 17, but rather map to mouse chromosome 10. Unlike the GPI-linked Rae-1 proteins, H60 appears to be anchored to the membrane via a hydrophobic transmembrane segment. H60 transcripts were found in embryonic tissue, in spleen, and in some transformed cell lines. Transcripts were also observed in mouse skin cells after exposure to carcinogens. Binding of H60 to NKG2D results in the activation of cytolytic activity and/or cytokine production by the NKG2D-expressing effector cells. Ectopic expression of H60 on mouse tumor cell lines resulted in the *in vivo* rejection of the tumors (1-6).

References:

1. Malarkannan, S. *et al.* (1998) *J. Immunol.* **161**:3501.
2. Diefenbach, A. *et al.* (2000) *Nature Immunol.* **1**:119.
3. Cerwenka, A. *et al.* (2000) *Immunity* **12**:721.
4. Cerwenka, A. *et al.* (2001) *Proc. Natl. Acad. Sci. USA* **98**:11521.
5. Diefenbach, A. *et al.* (2001) *Nature* **413**:165.
6. NKG2D and its Ligands, www.RnDSystems.com.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.