

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Integrin α V/CD51 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 273210
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Integrin α V/CD51 Phe31-Val992 Accession # NP_002201
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 μ g/10 ⁶ cells	Human peripheral blood mononuclear cells

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Integrin α V (CD51; also vitronectin receptor subunit alpha) is a 140-150 kDa member of the integrin α -chain family of adhesion molecules. It forms disulfide-linked integral membrane heterodimers with at least five β -chains, including β 1, 3, 5, 6 and 8. Human α V is a 1018 aa type I transmembrane glycoprotein that contains a 962 aa extracellular domain (ECD) and a short 32 aa cytoplasmic tail. The ECD contains seven FG (PheAlaGly)-GAP (GlyAlaPro) repeats that form a β -propeller domain (aa 46-483). Furin cleavage of the α V ECD occurs after Gly889, generating a disulfide-linked heteromeric subunit α V chain. α V-containing integrins bind multiple ECM molecules, including vitronectin, osteopontin, MMP-2 and TSP. The ECD of human α V shares 92% and 90% aa sequence identity with mouse and rat α V ECD, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.