

Human/Mouse/Rat/Chicken Oligodendrocyte Marker O4 APC-conjugated Antibody

Monoclonal Mouse IgM Clone # O4 Catalog Number: FAB1326A 100 TESTS


DESCRIPTION			
Species Reactivity	Human/Mouse/Rat/Chicken		
Specificity	Detects human, mouse, rat, and chicken Oligodendrocyte Marker O4.		
Source	Monoclonal Mouse IgM Clone # O4		
Purification	IgM-specific Affinity-purified from hybridoma culture supernatant		
Immunogen	Bovine brain corpus callosum white matter		
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of Oligodendrocyte Marker O4 in Differentiated Rat Cortical Stem Cells by Flow Cytometry.

Differentiated Rat Cortical Stem Cells (Catalog # NSC001) were stained with Mouse Anti-Human/Mouse/Rat/Chicken Oligodendrocyte Marker O4 APC-conjugated Monoclonal Antibody (Catalog # FAB1326A, filled histogram) or isotype control antibody (Catalog # IC015A, open histogram). View our protocol for Staining Membrane-associated Proteins

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Oligodendrocytes are myelinating cells in the central nervous system (CNS) and form the myelin sheath of axons to support rapid nerve conduction. Oligodendrocyte Marker O4 is an antigen on the surface of oligodendrocyte progenitors (1, 2). It has been commonly used as the earliest recognized marker specific for the oligodendroglial lineage (3-8).

References:

- 1. Schachner, M. et al. (1981) Dev. Biol. 83:328.
- 2. Bansal, R. et al. (1989) J. Neurosci. Res. 24:548.
- 3. Bansal, R. and Pfeiffer, S.E. (1989) Proc. Natl. Acad. Sci. USA 86:6181.
- 4. Gard, A. et al. (1995) Dev. Biol. 167:596.
- 5. Reynolds, R. and Hardy, R. (1997) J. Neurosci. Res. 47:455.
- 6. Ono, K. et al. (1997) J. Neurosci. Res. 48:212.
- 7. Pang, Y. et al. (2000) J. Neurosci. Res. 62:510.
- 8. Cai, Z. et al. (2001) Brain Res. 898:126.

