

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BCAM in ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human (rh) ALCAM, rhEpCAM, recombinant mouse (rm) MAdCAM-1, rhMCAM, rhNCAM-L1, rmOCAM, or rmTROP-2 is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 87207
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human BCAM Glu32-Ala547 Accession # CAA58449
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Huh-7 human hepatoma cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Basal-Cell Adhesion Molecule (BCAM) and Lutheran blood group glycoprotein (LU) are two alternatively spliced variants of a single immunoglobulin superfamily (IgSF) protein that differ in the length of their cytoplasmic tails. BCAM cDNA encodes a 628 amino acid (aa) residues precursor protein with a putative 31 aa signal peptide, a 597 aa extracellular domain containing three C2 type and two V-type Ig-like domains, a 21 aa transmembrane domain, and a 19 aa cytoplasmic domain. Compared to the 40 aa cytoplasmic domain present in LU, the BCAM cytoplasmic tail lacks the putative Src homology 3 (SH3) binding site that may be involved in mediating intracellular signaling. BCAM/LU has wide tissue distribution and is expressed on erythrocytes, the endothelium of blood vessels and on the basal layer of cells in the epithelia. The expression of BCAM/LU in normal tissues is higher in fetal versus adult tissues. BCAM/LU expression is also upregulated in sickle cell disease red blood cells, in activated keratinocytes and following malignant transformation in some cell types *in vivo* and *in vitro*. BCAM/LU has been shown to be an adhesion molecule that binds laminin, a basement membrane protein involved in cell differentiation, adhesion, migration and proliferation.

References:

- Campbell, I.G. *et al.* (1994) *Cancer Research* **54**:5761.
- Parsons, S.F. *et al.* (1995) *Proc. Natl., Acad. Sci. USA*, **92**:5496.
- Udani, M. *et al.* (1998) **101**:2550.
- Schon, M. *et al.* (2000) *J. Invest. Dermatol.*, **115**:1047.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.