

Human DLL4 PE-conjugated Antibody

Monoclonal Rat IgG_{2A} Clone # 447506

Catalog Number: FAB1506P 100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human DLL4 in direct ELISAs.		
Source	Monoclonal Rat IgG _{2A} Clone # 447506		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant human DLL4 Ser27-Pro524 Accession # Q9NR61		
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of DLL4 in HUVEC Human Cells by Flow Cytometry. HUVEC human umbilical vein endothelial cells were stained with Rat Anti-Human DLL4 PE-conjugated Monoclonal Antibody (Catalog # FAB1506P, filled histogram) or isotype control antibody (Catalog # IC006P, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Delta-like ligand 4 (DLL4) is a transmembrane protein that contains one DSL domain and eight tandem EGF-like repeats. DLL4 is expressed on arterial endothelial cells where it signals through Notch 1 and Notch 4. DLL4 expression is induced by VEGF and regulates the proliferation of endothelial tip cells during vascular sprouting. DLL4 blockade as well as overexpression can suppress tumor growth by promoting dysregulated angiogenesis. Within the extracellular domain, human DLL4 shares 85% amino acid sequence identity with mouse and rat DLL4.

