

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CCR-2 transfectants but not the parental cell line or CCR-5 transfectants.
Source	Monoclonal Mouse IgG _{2B} Clone # 48607
Purification	Protein A or G purified from ascites
Immunogen	NS0 mouse myeloma cell line transfected with human CCR2 Met1-Leu360 Accession # NP_001116868
Conjugate	PerCP (Peridinin-chlorophyll Protein Complex) Excitation Wavelength: 482 and 564 nm Emission Wavelength: 675 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CCR2 is a G-protein linked seven transmembrane domain spanning chemokine receptor that preferentially binds monocyte chemoattractant proteins-1 and -3 (MCP-1 and MCP-3). Two isoforms of this receptor (CCR2A and CCR2B) are expressed on cell surfaces as a result of alternate splicing from the same gene. These two CCR2 variants differ only at their intracellular carboxyl terminals, with the CCR2A form possessing 14 additional amino acids. This may provide a mechanism by which cells responding to similar extracellular ligands can activate different intracellular second messengers. Cells that respond to the action of MCP-1 and therefore are likely to express CCR2 receptors, include monocytes, T cells, NK cells, basophils, mast cells and dendritic cells. A recent report suggests that B cells may also express CCR2 receptors. The recognition that a variety of chemokine receptors, including CCR2, can serve as HIV fusion co-factors and as facilitators of T cell recruitment during inflammation makes chemokine receptor monitoring an important exercise in elucidating the HIV infection process and the regulation of inflammatory reactions.