

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse ICOS in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 670306
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with mouse ICOS Accession # Q9WVS0
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Mouse splenocytes treated with Concanavalin A

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Mouse ICOS (inducible co-stimulator), also called ALLIM (activation-inducible lymphocyte immunomediatory molecule) and CRP-1 (CD28-related protein-1), is a member of the CD28 family of immune costimulatory receptors. Other family members are CD28, CTLA-4 and PD-1. Mouse ICOS is a homodimeric type I transmembrane protein consisting of 200 amino acids (aa) with a putative 20 aa signal sequence, a 122 aa extracellular domain, a 23 aa transmembrane region, and a 35 aa cytoplasmic domain. ICOS shares approximately 39% aa similarity with CD 28 and CTLA-4. Mouse and human ICOS share approximately 72% aa identity. ICOS is expressed on most CD45RO⁺ cells. ICOS expression is upregulated within approximately 24-48 hours of activation on T_H primed cells. B7-H2, a member of the B7 family of costimulatory ligands, has been identified as the ICOS ligand. The B7-H2/ICOS interaction appears to play roles in T cell dependent B cell activation and T_H differentiation.

References:

1. Coyle, A.J. and J.C. Gutierrez-Ramos (2001) *Nat. Immunol.* **2**:203.
2. Mages, H.W. *et al.* (2000) *Eur. J. Immunol.* **30**:1040.
3. Yoshinaga, S.K. *et al.* (1999) *Nature* **402**:827.
4. Hutloff, A. *et al.* (1999) *Nature* **397**:263.
5. Aicher, A. *et al.* (2000) *J. Immunol.* **164**:4689.
6. Coyle, A.J. *et al.* (2000) *Immunity* **13**:95.
7. Gonzalo, J.A. *et al.* (2001) *J. Immunol.* **166**:1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.