

# Human Cadherin-11 PE-conjugated Antibody

Monoclonal Mouse IgG<sub>2A</sub> Clone # 667039

Catalog Number: FAB17901P 100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human Cadherin-11 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) Cadherin-4, -6, -8, -12, -13, -17, rhE-Cadherin, rhN-Cadherin, rhP-Cadherin, or rhVE-Cadherin is observed.		
Source	Monoclonal Mouse IgG <sub>2A</sub> Clone # 667039		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Cadherin-11 Phe23-Thr617 Accession # AAA35622		
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.		

### **APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 <sup>6</sup> cells	See Below

## DATA



Detection of Cadherin-11 in PC-3 Human Cell Line by Flow Cytometry. PC-3 human prostate cancer cell line was stained with Mouse Anti-Human Cadherin-11 PE-conjugated Monoclonal Antibody (Catalog # FAB17901P, filled histogram) or isotype control antibody (Catalog # IC003P, open histogram). Cells were stained in a buffer containing Ca<sup>2+</sup> and Mg<sup>2+</sup>. View our protocol for Staining Membrane-associated Proteins.

## PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.





## Human Cadherin-11 PE-conjugated Antibody

Monoclonal Mouse IgG<sub>2A</sub> Clone # 667039

Catalog Number: FAB17901P

100 TESTS

## BACKGROUND

Cadherin-11, also known as OB-Cadherin, is a 120 kDa member of the classical Cadherin family of calcium-dependent homophilic adhesion proteins. Cadherins are involved in multiple processes including embryonic development, cell migration, and maintenance of epithelial integrity (1). Cadherin-11 is expressed in embryonic mesodermal tissues and contributes to the morphogenesis of the nervous and skeletal systems (2-4). It is expressed on osteoblasts in the adult where it promotes the differentiation of both osteoblasts and chondrocytes (5). Cadherin-11 is up-regulated on breast cancer and prostate cancer cells which preferentially metastasize to bone (6, 7). It facilitates this metastasis via homophilic adhesion to bone marrow stroma and osteoblast-expressed Cadherin-11 (6-8). In the synovium, Cadherin-11 supports adhesion between synoviocytes but promotes cell invasion in synovitis and rheumatoid arthritis (9, 10). Its up-regulation in the vasculature following injury contributes to intimal hyperplasia by inducing smooth muscle cell migration and proliferation (11). In the nervous system, Cadherin-11 interacts with FGF R1 to promote neurite extension from spinal cord explants (12). Mature human Cadherin-11 consists of a 564 amino acid (aa) extracellular domain (ECD) with five tandem Cadherin repeats, a 23 aa transmembrane segment, and a 156 aa cytoplasmic domain (13, 14). Within the ECD, human Cadherin-11 shares 97% and 98% aa sequence identity with mouse and rat Cadherin-11, respectively. An 80 kDa portion of the Cadherin-11 ECD can be shed by proteolytic cleavage, and this fragment competes with the full length molecule for cell adhesion (3, 15). Alternate splicing of human Cadherin-11 generates an 85 kDa isoform with substituted transmembrane and cytoplasmic regions (14, 15).

#### References:

- 1. Pokutta, S. and W.I. Weis (2007) Annu. Rev. Cell Dev. Biol. 23:237.
- 2. Kimura, Y. et al. (1995) Dev. Biol. 169:347.
- 3. McCusker, C. et al. (2009) Mol. Biol. Cell 20:78.
- 4. Clendenon, S.G. et al. (2009) Dev. Dyn. 238:1909.
- 5. Kii, I. et al. (2004) J. Bone Mineral Res. 19:1840.
- 6. Tamura, D. et al. (2008) Int. J. Oncol. 33:17.
- 7. Chu, K. et al. (2008) Mol. Cancer Res. 6:1259.
- 8. Huang, C.-F. et al. (2010) Cancer Res. 70:4580.
- 9. Valencia, X. et al. (2004) J. Exp. Med. 200:1673.
- 10. Kiener, H.P. et al. (2009) Arthritis Rheum. **60**:1305.
- Monahan, T.S. et al. (2007) J. Vasc. Surg. 45:581.
   Boscher, C. and R.-M. Mege (2008) Cell. Signal. 20:1061.
- 13. Tanihara, H. et al. (1994) Cell Adhes. Commun. 2:15.
- 14. Okazaki, M. et al. (1994) J. Biol. Chem. 269:12092.
- 15. Kawaguchi, J. et al. (1999) J. Bone Mineral Res. 14:764.

