

Human CRACC/SLAMF7 Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG_{2A} Clone # 235614

Catalog Number: FAB1906G

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human CRACC/SLAMF7 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human NTB-AR is observed.		
Source	Monoclonal Mouse IgG _{2A} Clone # 235614		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CRACC/SLAMF7 Lys27-Ser225 Accession # Q9NQ25		
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website

	Recommended Concentration	Sample
Flow Cytometry	5 μL/10 ⁶ cells	See Below

DATA

Flow Cytometry



Detection of CRACC/SLAMF7 in Human PBMCs by Flow Cytometry. Human peripheral blood mononuclear cells (PBMCs) were stained with Mouse Anti-Human CD3c APC-conjugated Monoclonal Antibody (Catalog # FAB100A) and either (A) Mouse Anti-Human CRACC/SLAMF7 Alexa Fluor® 488-conjugated Monoclonal Antibody (Catalog # FAB1906G) or (B) Mouse IgG_{2A} Alexa Fluor 488 Isotype Control (Catalog # IC003G). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

ShippingThe product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CRACC, also known as CSI (CD2 subset I) and CD319, is a type I transmembrane protein belonging to the CD2 subset of the Ig superfamily. CRACC is expressed on most NK cells and subsets of CD8⁺ cells, CD4⁺ cells and B cells. CRACC may play a role in the activation and effector function of T cells and NK cells and likely participates in B cell adhesion. Over amino acids (aa) 27-225, human and mouse SLAMF7 share 53% as sequence identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 2/6/2018 Page 1 of 1

