

#### DESCRIPTION

<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human and mouse Frizzled-4 in Western blots. In Western blots, no cross-reactivity with recombinant mouse (rm) Frizzled-1, -2, -3, -6, -7, -8, or recombinant human Frizzled-5 is observed. Detects Frizzled-4 by flow cytometry and immunohistochemistry on mouse differentiated embryoid bodies derived from the mouse embryonic cell line, D3.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 145901
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse Frizzled-4 Phe37-Glu180 Accession # Q61088
<b>Conjugate</b>	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	0.25-1 µg/10 <sup>6</sup> cells	HEK293 human embryonic kidney cell line

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

#### BACKGROUND

Frizzled-4, designated CD344, is a 7-transmembrane glycoprotein of the Frizzled family within the G-protein coupled receptor superfamily (1, 2). Frizzled proteins function as receptors for Wnt proteins and can activate canonical Wnt/beta-catenin signaling as well as planar cell polarity and calcium flux pathways (1). Frizzled-4 is particularly important in angiogenic Wnt pathway signaling (1, 5). Frizzleds contain a divergent N-terminal signal peptide, a highly conserved ~120 amino acid (aa) cysteine-rich domain (CRD), a variable length linker region, seven transmembrane domains, and a variable-length C-terminal tail (1). The mouse Frizzled-4 cDNA encodes 537 aa with a 36 aa signal sequence and a 186 aa N-terminal extracellular sequence (4). The portion expressed includes aa 37-180, and shares 93%, 94%, 90%, 89% and 88% identity with human, rat, equine, bovine and canine Frizzled-4, respectively. This portion competes for Wnt binding with endogenous receptors. In humans, a 122 aa soluble form that diverges at aa 95 is proposed to be a positive regulator of Wnt signaling pathways (5). Frizzled-4 is unusual in binding a non-wnt ligand, Norrin, in addition to binding Wnt ligands (1, 3, 6). Norrin binds the Frizzled-4 CRD, activates Wnt signaling pathways and uses LRP5/6 as co-receptors (3, 6). Deletion of either Frizzled-4 or Norrin in mice results in a similar phenotype including malformation of vasculature in the retina, cerebellar degeneration, and loss of hair cells in the inner ear (1, 3, 7). In humans, blindness due to familial exudative vitreoretinopathy (FEVR) is associated with mutations producing loss of function of Frizzled-4 or Norrin, designated EVR1 and EVR2, respectively (1, 3, 8). Frizzled-4 expression has been found in many tissues, including mouse ovary, where it influences corpus luteum vasculogenesis and is necessary for fertility (4, 9).

#### References:

- Huang, H-C. and P.S. Klein (2004) *Genome Biol.* **5**:234.
- Parmalee, N.L. and J. Kitajewski (2008) *Curr. Drug Targets* **9**:558.
- Xu, Q. *et al.* (2004) *Cell* **116**:883.
- Wang, Y. *et al.* (1996) *J. Biol. Chem.* **271**:4468.
- Sagara, N. *et al.* (2001) *Biochem. Biophys. Res. Commun.* **282**:750.
- Smallwood, P.M. *et al.* (2007) *J. Biol. Chem.* **282**:4057.
- Wang, Y. *et al.* (2001) *J. Neurosci.* **21**:4761.
- Robitaille, J. *et al.* (2002) *Nat. Genet.* **32**:326.
- Hsieh, M. *et al.* (2005) *Biol. Reprod.* **73**:1135.

# Human/Mouse Frizzled-4 Alexa Fluor® 700-conjugated Antibody

Monoclonal Rat IgG<sub>2A</sub> Clone # 145901

Catalog Number: FAB194N  
100 µg

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.