

DESCRIPTION	
Species Reactivity	Mouse/Rat
Specificity	Detects mouse MIF in ELISA and Western Blot.
Source	Monoclonal Rat IgM Clone # 811409
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse MIF Met1-Ala115 Accession # P34884
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	J774A.1 mouse reticulum cell sarcoma macrophage cell line fixed with paraformaldehyde and permeabilized with saponin

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

MIF (Macrophage Inhibitory Factor; also GIF) is a 12-13 kDa member of the MIF family of proteins. Although secreted, it possesses no definitive signal sequence. MIF is widely expressed and serves to regulate immune reactions. On macrophages and B cells, MIF binds to a CD74:CD44 complex, initiating downstream signaling. On monocytes, this promotes TNFα production plus IFN-γ-stimulated NO production. On B cells, this promotes B cell survival. MIF also binds to leukocyte CD74:CXCR2 and CD74:CXCR4 heterodimers, initiating T cell and monocyte migration. A tautomerase activity has also been shown for MIF and is suggested to involve the entire length of the molecule. Mouse MIF is 114 amino acids (aa) in length. MIF is phosphorylated at Ser91, and an additional, singular Cys residue is covalently linked to MIF at Cys60. MIF is considered to act as a homotrimer. Full-length mouse MIF shows 90% and 99% aa sequence identity with human and rat MIF, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.