

Human/Mouse SSEA-1 PE-conjugated Antibody

Monoclonal Mouse IgM Clone # MC-480 Catalog Number: FAB2155P 100 TESTS


DESCRIPTION			
Species Reactivity	Human/Mouse		
Specificity	Detects human and mouse SSEA-1.		
Source	Monoclonal Mouse IgM Clone # MC-480		
Purification	IgM-specific Affinity-purified from hybridoma culture supernatant		
Immunogen	F9 mouse teratocarcinoma stem cells		
Conjugate	Phycoerythrin		
	Excitation Wavelength: 488 nm		
	Emission Wavelength: 565-605 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet		
	(SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Stage-Specific Embryonic Antigen-1 (SSEA-1), an antigenic epitope defined as a Lewis x carbohydrate structure, is expressed during early mouse embryogenesis on murine embryonal carcinoma cells (EC), murine embryonic stem cells (ES), and murine and human germ cells. Expression of SSEA-1 is down-regulated following differentiation of murine EC and ES cells. In contrast, the differentiation of human EC and ES cells is accompanied by an increase in SSEA-1 expression (1, 2).

References:

- 1. Solter, D. and Knowles, B.B. (1978) Proc. Natl. Acad. Sci. USA 75:5565.
- 2. Fox, N. et al. (1983) Cancer Res. 43:669.

