

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Cadherin-12 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) Cadherin-8, -11, -13, -17, rhE-Cadherin, rhP-Cadherin, rhR-Cadherin, or rhVR-Cadherin is observed.
Source	Monoclonal Rat IgG ₁ Clone # 343621
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	S. <i>frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human Cadherin-12 Met1-Pro609 Accession # P55289
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human neural progenitor cells differentiated by growth factor withdrawal

Note: Since classic Cadherins can be protected from trypsin treatment in the presence of Ca²⁺, cells in monolayer cultures are harvested with 0.01% Trypsin in the presence of 1-5 mM CaCl₂ at 37° C. Flow cytometry can be performed according to the standard procedures, except that all the cell staining and washing steps are performed in the presence of Ca²⁺ and Mg²⁺ (e.g. using FACS buffer: PBS containing 1 mM CaCl₂, 1 mM MgCl₂, 2% FBS and 0.02% sodium azide).

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

The cadherin superfamily is a large family of membrane-associated glycoproteins that engage in homotypic, calcium-dependent, cell-cell adhesion events. The superfamily can be divided into at least four subfamilies based on its member's extracellular (EC) regions and cytoplasmic domains (1, 2). These include classical cadherins, desmosomal cadherins, protocadherins, and cadherin-like molecules that contain a variable number of EC and transmembrane (TM) domains (1). Cadherin-12, also known as brain-cadherin and N-cadherin 2, is a 150 kDa classical cadherin. Classical family molecules are modular in their extracellular region, mediating calcium-dependent cell-cell adhesion through their five EC Ca²⁺-binding repeats (2). Cadherin-12 can be further identified as a type II classical cadherin, due to the absence of a His-Ala-Val motif in its most N-terminal cadherin repeat (3). Human Cadherin-12 is synthesized as a 794 amino acid (aa) type I transmembrane preproprotein that contains a 23 aa signal peptide, a 31 aa prosequence, a 555 aa extracellular region, a 28 aa transmembrane segment, and a 157 aa cytoplasmic domain (4, 5). The five EC cadherin domains are approximately 110 aa in length and generate two β-sheets that are oriented like bread in a sandwich. Human Cadherin-12 EC region is 96% aa identical to mouse Cadherin-12 EC region. Cadherin-12 is expressed specifically in CNS neurons. The bulk of its expression is postnatal, and it is proposed to be involved in synaptogenesis (4). As a classic cadherin, Cadherin-12 will form homodimers and promote intercellular adhesion with itself and, possibly, cadherins-8 and -14 (6).

References:

1. Koch, A.W. *et al.* (2004) *Cell. Mol. Life Sci.* **61**:1884.
2. Angst, B.D. *et al.* (2001) *J. Cell Sci.* **114**:629.
3. Gessner, R. and R. Tauber (2000) *Ann. N.Y. Acad. Sci.* **915**:136.
4. Selig, S. *et al.* (1997) *Proc. Natl. Acad. Sci. USA* **94**:2398.
5. Tanihara, H. *et al.* (1994) *Cell Adhes. Commun.* **2**:15.
6. Shimoyama, Y. *et al.* (2000) *Biochem. J.* **349**:159.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.