

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human 4-1BB Ligand/TNFSF9 Ligand in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant mouse (rm) 4-1BB Ligand, recombinant human (rh) APRIL, rhBAFF, rmEDA-A2, rhFas Ligand, rhGITR Ligand, rhLIGHT, rhLymphotoxin α1/β2, rhLymphotoxin α2/β1, rhOX40 Ligand, rhTNF-α, rhTRAIL, rhTRANCE, or rhTWEAK is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 282220
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human 4-1BB Ligand/TNFSF9 Arg71-Glu254 Accession # P41273.1
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Daudi human Burkitt's lymphoma cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

4-1BB Ligand (4-1BBL; also CD137L) is a 32 kDa type II transmembrane protein that belongs to the TNF superfamily (TNFSF) molecules (1-4). The human 4-1BBL cDNA encodes a 254 amino acid (aa) protein that contains a 25 aa N-terminal cytoplasmic domain, a 23 aa transmembrane segment, and a 206 aa C-terminal extracellular region (5). The extracellular domain (ECD) of 4-1BBL has a jelly-roll, β-sandwich tertiary structure that is similar to other TNFSF members. There is only one cysteine in the human ECD, and no potential N-linked glycosylation sites. The potential exists, however, for O-linked glycosylation. The human 4-1BBL ECD shares 32% and 35% aa identity with mouse and rat ECD, respectively. In the cytoplasmic domain, human 4-1BBL is 55 aa shorter than the equivalent region in rodents. 4-1BBL is expressed by activated B cells, macrophages, dendritic cells, activated T cells, neurons and astrocytes (2, 3, 6). A bioactive 26 kDa soluble form of 4-1BBL, presumably generated by MMP cleavage, occurs in humans (4). Human 4-1BBL signals through both CD137/4-1BB and itself. Its cytoplasmic tail participates in reverse signaling that induces apoptosis in T cells and cytokine secretion (IL-6; TNF-α) by monocytes (7, 8). 4-1BBL binding to CD137/4-1BB produces a number of effects. It seems to play a key role in the T cell recall response. It maintains T cell numbers at the end of a primary response, and induces CD4⁺ and CD8⁺ T cells to proliferate and secrete cytokines such as IL-2 and IFN-γ in CD4⁺ cells, and IFN-γ in CD8⁺ cells (9, 10).

References:

1. Kwon, B. *et al.* (2003) *Exp. Mol. Med.* **35**:8.
2. Salih, H.R. *et al.* (2002) *Int. J. Clin. Pharmacol. Ther.* **40**:348.
3. Vinay, D.S. and B.S. Kwon (1998) *Semin. Immunol.* **1**:481.
4. Salih, H.R. *et al.* (2001) *J. Immunol.* **167**:4059.
5. Alderson, M.R. *et al.* (1994) *Eur. J. Immunol.* **24**:2219.
6. Reali, C. *et al.* (2003) *J. Neurosci. Res.* **74**:67.
7. Michel, J. *et al.* (1999) *Immunology* **98**:42.
8. Langstein, J. *et al.* (1998) *J. Immunol.* **160**:2488.
9. Wen, T. *et al.* (2002) *J. Immunol.* **168**:4897.
10. Dawicki, W. and T.H. Watts (2004) *Eur. J. Immunol.* **34**:743.

Human 4-1BB Ligand/TNFSF9 Alexa Fluor® 647-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 282220

Catalog Number: FAB2295R

100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.