

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human NCAM-1/CD56 in ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) ALCAM, rhBCAM, rhEPCAM, rhMCAM, rhNCAM-1-L1, recombinant mouse (rm) MAdCAM-1, rmNCAM-1-L1, or rmOCAM is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 301040
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human NCAM-1/CD56
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Neural cell adhesion molecule 1 (NCAM-1) is a multifunctional member of the Ig superfamily. It belongs to a family of membrane-bound glycoproteins that are involved in Ca⁺⁺ independent cell matrix and homophilic or heterophilic cell-cell interactions. NCAM-1 specifically binds to heparan sulfate proteoglycans (1), the extracellular matrix protein agrin (2), and several chondroitin sulfate proteoglycans that include neurocan and phosphocan (3). There are three main forms of human NCAM-1 that arise by alternate splicing. These are designated NCAM-120/NCAM-1 (761 amino acids [aa]), NCAM-140 (848 aa), and NCAM-180 (1120 aa). NCAM-120 is GPI-linked, while NCAM-140 and NCAM-180 are type I transmembrane glycoproteins (4-6). Additional alternate splicing adds considerable diversity to all three forms, and extracellular proteolytic processing is possible for NCAM-180 (7-8). NCAM-1 is synthesized as a 761 aa preproprecursor that contains a 19 aa signal sequence, a 722 aa GPI-linked mature region, and a 20 aa C-terminal prosegment (4). The molecule contains five C-2 type Ig-like domains and two fibronectin type-III domains. Human to mouse, NCAM-1 is 93% aa identical. NCAM-1 appears to be highly sialylated. The polysialylation of NCAM-1 reduces its adhesive property and increases its neurite outgrowth promoting features (9). NCAM-1 in the adult brain shows a decline of sialylation relative to earlier developmental periods. In regions that retain a high degree of neuronal plasticity, however, the adult brain continues to express polysialylation-NCAM-1, suggesting sialylation of NCAM-1 is involved in regenerative processes and synaptic plasticity (10-13).

References:

1. Burg, M.A. *et al.* (1995) *J. Neurosci. Res.* **41**:49.
2. Storms, S.D. and U. Rutishauser (1998) *J Biol. Chem.* **273**:27124.
3. Margolis, R.K. *et al.* (1996) *Perspect. Dev. Neurobiol.* **3**:273.
4. Dickson, G. *et al.* (1987) *Cell* **50**:1119.
5. Lanier, L.L. *et al.* (1991) *J. Immunol.* **146**:4421.
6. Hemperly, J.J. *et al.* (1990) *J. Mol. Neurosci.* **2**:71.
7. Rutishauser, U. and C. Goridis (1986) *Trends Genet.* **2**:72.
8. Vawter, M.P. *et al.* (2001) *Exp. Neurol.* **172**:29.
9. Rutishauser, U. (1990) *Adv. Exp. Med. Biol.* **265**:179.
10. Becker, C.G. *et al.* (1996) *J. Neurosci. Res.* **45**:143.
11. Doherty, P. *et al.* (1995) *J. Neurobiol.* **26**:437.
12. Eckardt, M. *et al.* (2000) *J. Neurosci.* **20**:5234.
13. Muller, D. *et al.* (1996) *Neuron* **17**:413.