

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	This antibody has been validated by flow cytometry on TGF-β RII-transfected cells only. In contrast to Catalog # <a href="#">FAB2411A</a> , minimal staining of PBMC lymphocytes was observed with this antibody.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 25508
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human TGF-beta RII Ile24-Asp159 Accession # P37173.2
<b>Conjugate</b>	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 μL/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

TGF-β RII is a membrane bound serine/threonine kinase. Upon ligand binding, TGF-β RII interacts with TGF-β RI to form the heteromeric signaling complex that transduces TGF-β signals. A splice variant of the type II receptor, TGF-β RIIb, containing a 25 amino acid residue insertion near the N-terminus of the mature protein has also been described.