

DESCRIPTION


| | |
|---------------------------|--|
| Species Reactivity | Human |
| Specificity | Detects human LIF R α in direct ELISAs and Western blots. |
| Source | Monoclonal Mouse IgG ₁ Clone # 32953 |
| Purification | Protein A or G purified from ascites |
| Immunogen | <i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human LIF R α Gln45-Ser833 Accession # P42702 |
| Conjugate | Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm |
| Formulation | Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|----------------------------------|-----------|
| Flow Cytometry | 10 μ L/10 ⁶ cells | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

The activities of the pleiotropic cytokine LIF are mediated through a high-affinity heterodimeric receptor complex consisting of two membrane glycoproteins: an α subunit (LIF R α , also known as LIF R and CD118) that binds LIF with low affinity and the 130 kDa (gp130) subunit that does not bind LIF by itself, but is required for high-affinity binding of LIF by the complex. The gp130 subunit was first described as the signal transducing subunit of the high-affinity IL-6 receptor complex. Besides LIF, the high-affinity heterodimeric LIF receptor complex has been shown to mediate the activities of oncostatin M (OSM), cardiotrophin-1 and ciliary neurotrophic factor (CNTF).

Human LIF R α cDNA encodes a 1097 amino acid (aa) residue precursor type I membrane protein with a 44 aa residue signal peptide, a 789 aa residue extracellular domain, a 26 aa residue transmembrane domain, and a 238 aa residue cytoplasmic domain. LIF R α is a member of the cytokine receptor family and has extensive homology to gp130. The extracellular domain of LIF R α has two cytokine receptor domains and three fibronectin type III repeats. In mouse, mRNAs encoding a soluble LIF R α and lacking transmembrane and intracellular domains, have been isolated. Soluble LIF R α has been shown to bind LIF and has LIF antagonistic activity.

References:

1. Bazan, J.F. 1990, Proc. Natl. Acad. Sci. USA **87**:6934.
2. Gearing, D.P. (1994) *Guidebook to Cytokines and Their Receptors*, Academic Press, p130.
3. Pennica D. *et al.* (1995) J. Biol. Chem. **270**:10915.