

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Mouse |
| Specificity | Detects mouse CD36/SR-B3 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human CD36 is observed. |
| Source | Monoclonal Rat IgG ₁ Clone # 324205 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Chinese hamster ovary cell line CHO-derived recombinant mouse CD36/SR-B3 Gly30-Lys439 Accession # Q08857 |
| Conjugate | Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|---------------------------------|---|
| Flow Cytometry | 0.25-1 µg/10 ⁶ cells | J774A.1 mouse reticulum cell sarcoma macrophage cell line |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

CD36 (alternatively known as platelet membrane glycoprotein IV (GPIV), thrombospondin receptor, fatty acid translocase (FAT), and scavenger receptor class B, member 3 (SR-B3)) is an 88 kDa, integral membrane glycoprotein that belongs to the class B scavenger receptor family (1, 2). The molecule is described as being ditopic, with two transmembrane segments connected by an extracellular loop (3). Mouse CD36 is synthesized as a 472 amino acid (aa) protein that contains a 6 aa N-terminal cytoplasmic domain, a 22 aa N-terminal transmembrane segment, a 420 aa extracellular "loop", a 22 aa C-terminal transmembrane segment, and a 9 aa C-terminal cytoplasmic tail (4). Both cytoplasmic tails are palmitoylated, with the C-terminal tail involved in oxidized LDL binding (5, 6). With respect to the extracellular loop, the N-terminal region is believed to bind both thrombospondin-1 and Plasmodium-infected erythrocytes. Other ligands for CD36 include long-chain fatty acids, collagen, phospholipids and apoptotic cells (1). The extracellular loop of mouse CD36 shares 94%, 92%, 84% and 84% aa sequence identity with the extracellular loops of rat, hamster, human and bovine CD36, respectively. Cells known to express CD36 include capillary endothelium, adipocytes, skeletal muscle cells, intestinal epithelium, smooth muscle cells and hematopoietic cells such as RBC's, platelets and monocytes (1). On the surface of cells, CD36 is suggested to exist as a dimer in response to ligation (7). CD36 is reported to regulate fatty uptake, act as an angiogenic with TSP-1, and participate in the clearance of apoptotic phagocytes (1, 8).

References:

1. Febbraio, M. *et al.* (2001) *J. Clin. Invest.* **108**:795.
2. Silverstein, R.L. and M. Febbraio (2000) *Curr. Opin. Lipid.* **11**:483.
3. Gruarin, P. *et al.* (2000) *Biochem. Biophys. Res. Commun.* **275**:446.
4. Endemann, G. *et al.* (1993) *J. Biol. Chem.* **268**:11811.
5. Malaud, E. *et al.* (2002) *Biochem. J.* **364**:507.
6. Tao, N. *et al.* (1996) *J. Biol. Chem.* **271**:22315.
7. Daviet, L. *et al.* (1997) *Thromb. Haemost.* **78**:897.
8. Simantov, R. and R.L. Silverstein (2003) *Front. Biosci.* **8**:s874.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.