

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IL-5 R $\alpha$ /CD125 in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody does not cross-react with recombinant human (rh) IL-5 R $\beta$ , rhIL-9 R, rhIL-4 R, rhIL-5 R $\alpha$ , or rhIL-13 R $\alpha$ 1.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 26815
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human IL-5 R $\alpha$ /CD125 Asp21-Arg335
<b>Conjugate</b>	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Interleukin 5, produced primarily by activated T cells and mast cells, has diverse biological effects on a variety of cell types. Human IL-5 is a potent eosinophil differentiation and activation factor *in vivo* and *in vitro*. Additionally, it has also been reported that IL-5 can stimulate the proliferation and/or differentiation of basophils and B cells. The multiple effects of IL-5 are mediated by binding of the cytokine to specific cell surface receptors expressed on target cells. As is the case with many other cytokines, the functional high-affinity receptor for IL-5 is a complex consisting of a ligand binding subunit ( $\alpha$  chain) and a second subunit ( $\beta$  chain) that can modulate the ligand binding affinity of the receptor complex. In the case of IL-5, the  $\beta$  subunit is shared with the high affinity receptor complexes for IL-3 and GM-CSF. The  $\beta$  chain does not bind any of the cytokines in question but is indispensable for the cytokine-mediated signaling. cDNA clones for the  $\alpha$  chain (IL-5 R $\alpha$ ) of both the mouse and human high affinity IL-5 receptor complexes have been isolated. Human and mouse IL-5 R $\alpha$  are both members of the hematopoietin receptor superfamily characterized by the presence of the WSXWS, and a four cysteine residue motif in the extracellular domain of the transmembrane protein. In addition to the cDNA clone encoding the full-length transmembrane protein, cDNA clones that arise from alternative splicing and that encode soluble secreted forms of IL-5 R $\alpha$  have been isolated from mouse as well as human cells. A naturally-occurring soluble form of the IL-5 R $\alpha$  has been detected in biological fluids of autoimmune-prone mice and mice bearing chronic B cell leukemia (BCL<sub>1</sub>). A recombinant human IL-5 soluble receptor  $\alpha$  has been shown to bind the human IL-5 dimer in a 1:1 ratio and acts as a human IL-5 antagonist. This molecule inhibits the proliferation of IL-5-dependent cell lines and blocks human umbilical cord blood eosinophil differentiation.