
Human LAIR1 APC-conjugated Antibody Monoclonal Mouse IgG_{2B} Clone # 342219



Catalog Number: FAB2664A

100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human LAIR1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human LAIR2 or recombinant mouse LAIR1 is observed.		
Source	Monoclonal Mouse IgG _{2B} Clone # 342219		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LAIR1 isoform 1 Gln22-His163 Accession # Q6GTX8		
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of LAIR1 in Human Blood Lymphocytes by Flow Cytometry. Human peripheral blood lymphocytes were stained with Mouse Anti-Human LAIR1 APCconjugated Monoclonal Antibody (Catalog # FAB2664A, filled histogram) or isotype control antibody (Catalog # IC0041A, open histogram). View our protocol for Staining Membrane-associated Proteins

PREPARATION AND STORAGE

The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. Shipping

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

LAIR1 is an inhibitory receptor belonging to the Ig superfamily. It is a type I transmembrane protein with one extracellular Ig-like domain and two cytoplasmic ITIMs. Four LAIR1 splice variants exist. LAIR1b has a 17 aa deletion outside the Ig loop in the extracellular domain. It differs from LAIR1c by one aa residue. LAIR1d has a 77 aa truncation in the cytoplasmic domain. LAIR1 is expressed on NK cells, T cells, B cells, monocytes, dendritic cells and most thymocytes. The extracellular domain of human LAIR1 shares 40% aa identity with that of the mouse protein.

