

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Integrin α V β 3.
Source	Monoclonal Mouse IgG ₁ Clone # 23C6
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human osteoclasts
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	5 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Integrin α V β 3 together with α IIb β ₃, constitutes the only known β 3 Integrins (1-3). The non-covalent heterodimer of 170 kDa α V/CD51 and 93 kDa β ₃/CD61 subunits shows wide expression, notably by endothelial cells and osteoclasts (2-4). Each subunit has a transmembrane sequence and a short cytoplasmic tail connected to the cytoskeleton. Active cell surface α V β 3 adheres to matrix proteins including vitronectin, fibronectin, fibrinogen and thrombospondin (2, 3). The ligand binding site of α V β 3 is in the N-terminal head region, formed by interaction of the β 3 vWFA domain with the α V beta-propeller structure (4). The α V subunit contributes a thigh and a calf region, while the β 3 subunit contains a PSI domain and four cysteine-rich I-EGF folds. The α V subunit domains termed thigh, calf-1 and calf-2 generate a "knee" region that is bent when the α V β 3 is in its constitutively inactive state. Activation, either by "inside out" signaling or by Mg²⁺ or Mn²⁺ binding, extends the Integrin to expose its ligand binding site (1, 4). Two splice variants of β 3 (b and c) diverge over the last 21 amino acids (aa) and lack cytoplasmic phosphorylation sites (5, 6). Another β 3 splice variant diverges after the vWFA domain, producing a soluble 60 kDa form in platelets and endothelial cells (7). α V β 3 is essential for the maturation of osteoclasts and their binding and resorption of bone; it also, however, promotes their apoptosis (8, 9). M-CSF R and α V β 3 share signaling pathways during osteoclastogenesis, and deletion of either molecule causes osteopetrosis (8, 9). Also cell entry of several viruses is mediated by α V β 3 (4, 10). The 962 aa human α V ECD (11) shares 92-95% aa sequence identity with mouse, rat and cow α V while the 685 aa human β ₃ ECD (12) shares 95% aa identity with horse and dog, and 89-92% aa identity with mouse, rat and pig β ₃.

References:

1. Hynes, R. O. (2002) *Cell* **110**:673.
2. Serini, G. *et al.* (2006) *Exp. Cell Res.* **312**:651.
3. Ross, F. P. and S. L. Teitelbaum (2005) *Immunol. Rev.* **208**:88.
4. Xiong, J. *et al.* (2001) *Science* **294**:339.
5. Kumar, C. S. *et al.* (1987) *J. Biol. Chem.* **272**:16390.
6. vanKuppevelt, H. *et al.* (1989) *Proc. Natl. Acad. Sci. USA* **86**:5415.
7. Djaffar, I. *et al.* (1994) *Biochem. J.* **300**:67.
8. McHugh, K. P. *et al.* (2000) *J. Clin. Invest.* **105**:433.
9. Faccio, R. *et al.* (2003) *J. Clin. Invest.* **111**:749.
10. Chu, J. J. and M. Ng (2004) *J. Biol. Chem.* **279**:54533.
11. Suzuki, S. *et al.* (1987) *J. Biol. Chem.* **262**:14060.
12. Fitzgerald, L. A. *et al.* (1987) *J. Biol. Chem.* **262**:3936.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose, Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.