

# Human LINGO-1 APC-conjugated Antibody

Monoclonal Mouse IgG<sub>1</sub> Clone # 332237

Catalog Number: FAB30861A 100 TESTS


DESCRIPTION		
Species Reactivity	Human	
Specificity	Detects human LINGO-1 in direct ELISA and Western blot. In direct ELISA and Western blot, approximately 10% cross-reactivity with recombinant human (rh) LINGO-2 is observed and no cross-reactivity with rhLINGO-3 is observed.	
Source	Monoclonal Mouse IgG <sub>1</sub> Clone # 332237	
Purification	Protein A or G purified from hybridoma culture supernatant	
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LINGO-1 Thr40-Thr556 Accession # NP_116197	
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm	
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

#### **APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 <sup>6</sup> cells	See Below

### DATA



## PREPARATION AND STORAGE

**Shipping**The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

LINGO-1 (LRRN6A) is a 614 amino acid (aa) transmembrane protein of the leucine-rich repeat (LRR) family, ribonuclease inhibitor subfamily. The four known LINGO proteins contain LRR and IgCAM domains in the extracellular portion and share 44–61% as sequence identity. LINGO-1 is restricted to the nervous system and is concentrated in the brain as a component of the NgR1/p75 and NgR1/Taj (TROY) signaling complexes. LINGO-1 negatively regulates neurite outgrowth and myelination. LINGO-1 is highly conserved, showing 99% as sequence identity between human, mouse and rat.

