

Human VEGF R1/FIt-1 APC-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 49560

Catalog Number: FAB321A

100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human VEGF R1/Flt-1 in direct ELISAs and Western blots. In Western blots, this antibody does not cross-react with recombinant mouse VEGF R1, recombinant human (rh) VEGF R2, rhVEGF R3, or rhVEGF R4.		
Source	Monoclonal Mouse IgG ₁ Clone # 49560		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	S. frugiperda insect ovarian cell line Sf 21-derived recombinant human VEGF R1/Flt-1		
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of VEGF R1/Flt-1 in HUVEC Human Cells by Flow Cytometry. HUVEC human umbilical vein endothelial cells were stained with Mouse Anti-Human VEGF R1/Flt-1 APC-conjugated Monoclonal Antibody (Catalog # FAB321A, filled histogram) or isotype control antibody (Catalog # IC002A, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Vascular Endothelial Growth Factor Receptor 1 (VEGF R1) is a receptor tyrosine kinase that is expressed primarily on endothelial cells and plays a role in vasculogenesis and angiogenesis. A soluble variant of VEGF R1 was also reported to bind VEGF and PIGF with high affinity and function as a potent VEGF antagonist.

