

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD117/c-kit in ELISAs and Western blots. In sandwich immunoassays and Western blots, no cross-reactivity with recombinant mouse CD117 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 47233
Purification	Protein A or G purified from ascites
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human CD117/c-kit Gln26-Thr520 Accession # P10721
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	TF-1 human erythroleukemic cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Stem cell factor receptor (CD117, the gene product of the *c-kit* proto-oncogene) and its ligand, stem cell factor (also named c-kit ligand, mast cell growth factor), play essential roles in gametogenesis, melanogenesis, and hematopoiesis. The human stem cell factor receptor cDNA encodes a 972 amino acid (aa) residue precursor membrane protein with a 25 aa residue signal peptide (experimentally determined), a 495 aa residue extracellular domain, a 23 aa residue transmembrane segment, and a 429 aa residue cytoplasmic domain. Stem cell factor receptor is a member of the type III subfamily of receptor tyrosine kinases (RTK) that also includes the receptors for M-CSF, Flt-3, PDGF, and VEGF. All class III RTKs are characterized by the presence of five immunoglobulin-like domains in their extracellular region and a split kinase domain in their intracellular region. SCF binding induces receptor homodimerization and signal transduction. SCF receptor is expressed in hematopoietic progenitor cells, normal B and T lymphocyte progenitor cells, mast cells, germ cells, melanocytes, neurons, glial cells, placenta, kidney, lung, and gut. In addition, SCF receptor expression has also been reported in a number of human tumor cell lines. SCF receptor can be proteolytically cleaved from the cell surface and high levels of soluble SCF receptor has been detected in cell conditioned medium and human plasma. Recombinant soluble SCF receptor binds SCF with high affinity and is a potent SCF antagonist.

References:

1. Broudy, V. (1997) *Blood* **90**:1345.
2. Vliagoftis, H. *et al.* (1997) *J. Allergy Clin. Immunol.* **100**:435.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.