

#### DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human AMICA/JAML in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse AMICA is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 401901
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human AMICA/JAML Leu20-Leu275 Accession # Q86YT9
<b>Conjugate</b>	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25-1 µg/10 <sup>6</sup> cells	Human whole blood monocytes

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

#### BACKGROUND

AMICA (adhesion molecule, interacting with CXADR antigen 1), also known as JAML, is a 65 kDa, heavily glycosylated transmembrane protein that belongs to the junctional adhesion molecule (JAM) subset of the immunoglobulin superfamily (1). JAM family molecules contribute to intercellular connections within epithelial and endothelial cell layers, and mediate their interactions with various hemopoietic cells (1). The human AMICA cDNA encodes a 384 amino acid (aa) precursor that includes a 19 aa signal sequence, a 256 aa extracellular domain (ECD) with two Ig-like domains, a 21 aa transmembrane segment, and a 98 aa cytoplasmic domain (2). Alternative splicing may generate isoforms with N- and C-terminal deletions. In contrast to other JAM family proteins, AMICA does not contain a cytoplasmic PDZ-binding motif (3). Within the ECD, human AMICA shares 58% and 63% aa sequence identity with mouse and rat AMICA, respectively. It shares 18%-20% aa sequence identity with the ECDs of human JAM-A, -B, -C, and JAM4. AMICA is expressed on the surface of granulocytes and monocytes and is upregulated during the differentiation of myeloid leukemia cells (2, 3). A motif in the ECD, which promotes dimerization of other JAM family proteins, is required for surface localization of AMICA (2). AMICA mediates the adhesion of monocytes to endothelial cells (2) and neutrophil migration across epithelial cell monolayers (3). This latter function involves specific interactions of AMICA with the coxsackie virus and adenovirus receptor (CXADR) in epithelial tight junctions (3). In particular, the membrane proximal Ig-like domain of AMICA binds the membrane-distal Ig-like domain of CXADR (3). AMICA does not appear to interact homophilically, as neutrophils adhere to immobilized CXADR but not to immobilized AMICA (3).

#### References:

1. Mandell, K.J. and C.A. Parkos (2005) *Adv. Drug Deliv. Rev.* **57**:857.
2. Moog-Lutz, C. *et al.* (2003) *Blood* **102**:3371.
3. Zen, K. *et al.* (2005) *Mol. Biol. Cell* **16**:2694.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.