

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ErbB3/Her3 in ELISAs. Does not cross-react with recombinant EGF R.
Source	Monoclonal Mouse IgG ₁ Clone # 66223
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human ErbB3/Her3 Ser20-Thr643 Accession # P21860
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	MCF-7 human breast cancer cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

ErbB3, also called Her3 (human epidermal growth factor receptor 3), is a type I membrane glycoprotein that is a member of the ErbB family of tyrosine kinase receptors. ErbB family members serve as receptors for the epidermal growth factor (EGF) family of growth factors. Among ErbB family members, ErbB3 is unique in that it contains a defective kinase domain. ErbB3 is expressed in keratinocytes, melanocytes, skeletal muscle cells, embryonic myoblasts and Schwann cells. Monomeric ErbB3 serves as a low affinity receptor for the heregulins (HRG). ErbB3 heterodimerizes with ErbB2 to form a high affinity receptor complex. In contrast, ErbB3 homodimerization or heterodimerization with ErbB4 forms a low affinity heregulin-binding complex. Because ErbB3 contains a defective kinase domain, the kinase domain of ErbB2 is responsible for initiating the tyrosine phosphorylation signal through the heterodimeric receptor. It has been found that a discrete three amino acid signal in the ErbB3 cytoplasmic domain is critical for transactivation of ErbB2. The cytoplasmic domain of ErbB3 also contains six consensus binding motifs for the SH2 domain of the regulatory p85 subunit of phosphoinositide 3-kinase (PI 3-kinase, PI3K) as well as one proline-rich consensus binding motif for the SH3 domain of p85. Human ErbB3 consists of 1342 amino acids (aa) with a 19 aa signal sequence, a 624 aa extracellular domain, a 21 aa transmembrane region, and a 678 aa cytoplasmic domain. ErbB3 appears to play roles in development, cancer, communication at the neuromuscular junction, and regulation of cell growth and differentiation.

References:

1. Kraus, M.H. *et al.* (1989) *Proc. Natl. Acad. Sci. USA* **86**:9193.
2. Plowman, G.D. *et al.* (1990) *Proc. Natl. Acad. Sci. USA* **87**:4905.
3. Carraway, K.L. 3rd *et al.* (1994) *J. Biol. Chem.* **269**:14303.
4. Emkey, R. and C.R. Kahn (1997) *J. Biol. Chem.* **272**:31172.
5. Sundaresan, S. *et al.* (1998) *Endocrinology* **139**:4756.
6. Hellyer, N.J. *et al.* (1998) *Biochem. J.* **333**:757.
7. Schaefer, G. *et al.* (1999) *J. Biol. Chem.* **274**:859.
8. Hellyer, N.J. *et al.* (2001) *J. Biol. Chem.* **276**:42153.
9. Schlessinger, J. (2000) *Cell* **103**:211.
10. Daly, R.J. (1999) *Growth Factors* **16**:255.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.