

Human CD31/PECAM-1 PE-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 9G11

Catalog Number: FAB3567P

100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human CD31/PECAM-1 in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human		
	(rh) E-Selectin, rhICAM-1, -2, -3, rhVCAM-1, or recombinant mouse VCAM-1 is observed.		
Source	Monoclonal Mouse IgG ₁ Clone # 9G11		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Activated HUVEC human umbilical vein endothelial cells		
Conjugate	Phycoerythrin		
	Excitation Wavelength: 488 nm		
	Emission Wavelength: 565-605 nm		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains < 0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data She		
	(SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of CD31/PECAM-1 in Human Granulocytes by Flow Cytometry. Human peripheral blood granulocytes were stained with Mouse Anti-Human CD31/PECAM-1 PE-conjugated Monoclonal Antibody (Catalog # FAB3567P, filled histogram) or isotype control antibody (Catalog # IC002P, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

The CD31 adhesion molecule, also known as PECAM-1, is expressed in large amounts on endothelial cells at intercellular junctions and on T cell subsets, and to a lesser extent on platelets and most other leukocytes such as monocytes and neutrophils. CD31 binds to itself homotypically, and also to the leukocyte integrin αν β3 heterotypically. CD31 is required for the transendothelial migration of leukocytes through intercellular junctions of vascular endothelial cells. CD31 has been found in human plasma, and the presence of this circulating isoform is suggested to modulate the transendothelial migration of leukocytes.

