

Human HGF R/c-MET Alexa Fluor® 594-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 95106

Catalog Number: FAB3582T
100 µg

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human HGF R/c-MET.
Source	Monoclonal Mouse IgG ₁ Clone # 95106
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human HGF R/c-MET Glu25-Thr932 Accession # P08581
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	MDA-MB-231 human breast cancer cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

HGF R, also known as Met (from *N*-methyl-*N*-nitro-*N*-nitrosoguanidine induced), is a glycosylated receptor tyrosine kinase that plays a central role in epithelial morphogenesis and cancer development. HGF R is synthesized as a single chain precursor which undergoes cotranslational proteolytic cleavage. This generates a mature HGF R that is a disulfide-linked dimer composed of a 50 kDa extracellular α chain and a 145 kDa transmembrane β chain (1, 2). The extracellular domain (ECD) contains a seven bladed β -propeller sema domain, a cysteine-rich PSI/MRS, and four Ig-like E-set domains, while the cytoplasmic region includes the tyrosine kinase domain (3, 4). Proteolysis and alternate splicing generate additional forms of human HGF R which either lack of the kinase domain, consist of secreted extracellular domains, or are deficient in proteolytic separation of the α and β chains (5-7). The sema domain, which is formed by both the α and β chains of HGF R, mediates both ligand binding and receptor dimerization (3, 8). Ligand-induced tyrosine phosphorylation in the cytoplasmic region activates the kinase domain and provides docking sites for multiple SH2-containing molecules (9, 10). HGF stimulation induces HGF R downregulation *via* internalization and proteasome-dependent degradation (11). In the absence of ligand, HGF R forms non-covalent complexes with a variety of membrane proteins including CD44v6, CD151, EGF R, Fas, Integrin $\alpha 6/\beta 4$, Plexins B1, 2, 3, and MSP R/Ron (12-19). Ligation of one complex component triggers activation of the other, followed by cooperative signaling effects (12-19). Formation of some of these heteromeric complexes is a requirement for epithelial cell morphogenesis and tumor cell invasion (12, 16, 17). Paracrine induction of epithelial cell scattering and branching tubulogenesis results from the stimulation of HGF R on undifferentiated epithelium by HGF released from neighboring mesenchymal cells (20). Genetic polymorphisms, chromosomal translocation, over-expression, and additional splicing and proteolytic cleavage of HGF R have been described in a wide range of cancers (1). Within the ECD, human HGF R shares 86-88% amino acid sequence identity with canine, mouse, and rat HGF R.

References:

1. Birchmeier, C. *et al.* (2003) *Nat. Rev. Mol. Cell Biol.* **4**:915.
2. Corso, S. *et al.* (2005) *Trends Mol. Med.* **11**:284.
3. Gherardi, E. *et al.* (2003) *Proc. Natl. Acad. Sci. USA* **100**:12039.
4. Park, M. *et al.* (1987) *Proc. Natl. Acad. Sci. USA* **84**:6379.
5. Crepaldi, T. *et al.* (1994) *J. Biol. Chem.* **269**:1750.
6. Prat, M. *et al.* (1991) *Mol. Cell. Biol.* **12**:5954.
7. Rodrigues, G.A. *et al.* (1991) *Mol. Cell. Biol.* **11**:2962.
8. Kong-Beltran, M. *et al.* (2004) *Cancer Cell* **6**:75.
9. Naldini, L. *et al.* (1991) *Mol. Cell. Biol.* **11**:1793.
10. Ponzetto, C. *et al.* (1994) *Cell* **77**:261.
11. Jeffers, M. *et al.* (1997) *Mol. Cell. Biol.* **17**:799.
12. Orian-Rousseau, V. *et al.* (2002) *Genes Dev.* **16**:3074.
13. Klosek, S.K. *et al.* (2005) *Biochem. Biophys. Res. Commun.* **336**:408.
14. Jo, M. *et al.* (2000) *J. Biol. Chem.* **275**:8806.
15. Wang, X. *et al.* (2002) *Mol. Cell* **9**:411.
16. Trusolino, L. *et al.* (2001) *Cell* **107**:643.
17. Giordano, S. *et al.* (2002) *Nat. Cell Biol.* **4**:720.
18. Conrotto, P. *et al.* (2004) *Oncogene* **23**:5131.
19. Follenzi, A. *et al.* (2000) *Oncogene* **19**:3041.
20. Sonnenberg, E. *et al.* (1993) *J. Cell Biol.* **123**:223.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.