

Human Integrin αL/CD11a Fluorescein-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 345913

Catalog Number: FAB3595F

100 TESTS


DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human Integrin αL/CD11a in Western blots.		
Source	Monoclonal Mouse IgG ₁ Clone # 345913		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Integrin αL/CD11a Tyr26-Met1089 (Tyr660lle) Accession # CAA68747		
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of Integrin $\alpha L/CD11a$ in Human Blood Lymphocytes by Flow Cytometry. Human peripheral blood lymphocytes were stained with Mouse Anti-Human Integrin $\alpha L/CD11a$ Fluorescein-conjugated Monoclonal Antibody (Catalog # FAB3595F, filled histogram) or isotype control antibody (Catalog # IC002F, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Integrin subunit αL/CD11a is a 180 kDa type I TM glycoprotein that interacts only with Integrin β2/CD18 to form LFA-1, a leukocyte adhesion protein which binds endothelial cell ICAM-1 through -4. Human Integrin αL contains a 1064 aa extracellular domain (ECD), a 20 aa TM sequence and a 58 aa cytoplasmic domain. The ECD contains seven repeats that form a beta-propeller structure and one inserted vWA domain (I domain) containing a metal ion-dependent adhesion site (MIDAS). Human and mouse Integrin αL ECD share 74% aa identity. CD11a has two splice variants, one which contains a 53 aa insert after Gln954, and a second which shows a deletion of aa 110-192.

