

Human Complement Component C5a R1 APC-conjugated Antibody

Monoclonal Mouse IgG_{2A} Clone # 347214

Catalog Number: FAB3648A 100 TESTS


DESCRIPTION		
Species Reactivity	Human	
Specificity	Detects human Complement Component C5a R1. Stains human Complement Component C5a R1 transfected cells but not irrelevant transfectants.	
Source	Monoclonal Mouse IgG _{2A} Clone # 347214	
Purification	Protein A or G purified from hybridoma culture supernatant	
Immunogen	NS0 mouse myeloma cell line transfected with human Complement Component C5a R1 Met1-Val350 Accession # P21730	
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm	
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



Detection of Complement
Component C5a R1 in Human
Granulocytes by Flow Cytometry. Human
granulocytes were stained with Mouse AntiHuman Complement Component C5a R1
APC-conjugated Monoclonal Antibody
(Catalog # FAB3648A, filled histogram) or
isotype control antibody (Catalog # IC003A,
open histogram). View our protocol for
Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

• 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Complement Component C5a R1 (C5aR1), also known as C5a ligand and CD88, is a 7TM protein expressed on myeloid, endothelial, epithelial, and smooth muscle cells. C5aR binds the activated complement anaphylatoxin C5a. In established allergic environments, this triggers neutrophil and eosinophil chemotaxis and the release of proinflammatory mediators. In contrast, C5aR1/C5a interactions are protective during allergen sensitization. Human C5aR1 shares 66% amino acid sequence identity with mouse and rat C5aR.

Rev. 10/14/2015 Page 1 of 1

