

## DESCRIPTION


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	In direct ELISAs, approximately 20% cross-reactivity with recombinant mouse (rm) CD97 and no cross-reactivity with recombinant human (rh) CD96v2, rhCD97, or rmCD96 is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>1</sub> Clone # 587702
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant mouse CD97v2 Gln24-His384 Accession # AAH06676
<b>Conjugate</b>	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	10 μL/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

CD97 is an approximately 100 kDa N-glycosylated member of the LNB-TM7 family of G protein-coupled proteins. It contains a 510 amino acid (aa) N-terminal extracellular region with four EGF-like domains and a juxtamembrane GPS motif which enables proteolytic shedding of the N-terminal extracellular domain. The alternately spliced variant 2 of CD97 exhibits a 94 aa deletion that results in loss of the third EGF-like repeat. CD97 is expressed on monocytes, macrophages, T cells, B cell subsets, dendritic cells, hematopoietic progenitor cells, and smooth muscle cells. It binds CD55/DAF, chondroitin sulfate, and Integrin although interaction with these molecules can vary between splice forms of CD97. CD97 participates in neutrophil migration, host defense, and angiogenesis. Within aa 24-384, mouse CD97 variant 2 shares 51% aa and 76% aa sequence identity with corresponding regions of human and rat CD97, respectively.