


DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human CD4 in direct ELISAs and Western blots. Does not cross-react with recombinant mouse CD4.
Source	Monoclonal Mouse IgG _{2A} Clone # 11830
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Recombinant human CD4 Extracellular domain Accession # P01730
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	5 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CD4 is a 54-60 kDa type I transmembrane glycoprotein that belongs to the immunoglobulin superfamily. It is expressed predominantly on hematopoietic cells such as T cells, monocytes and Langerhans cells. T cells known to be positive for CD4 include Th1, Th2, Th17, Th22, Th9, Tfh and Treg subsets, plus select thymocytes. CD4 exists as either an oxidized monomer (3 internal disulfide bonds), a reduced monomer (2 internal disulfide bonds) and a disulfide-linked dimer. CD4 functions in collaboration with the T cell receptor in the recognition of peptide antigens that are presented by class II major histocompatibility complexes. In particular, CD4 in its dimeric form binds to the MHC complex at non-polymorphic sites. Monomeric CD4 has also been shown to be a coreceptor for HIV entry and specifically to bind gp120, the external envelope glycoprotein of HIV. Over amino acids (aa) 26-396 (the extracellular domain), human and mouse CD4 share 58% aa sequence identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.