

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse M-CSF R/CD115 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human M-CSF R/CD115, recombinant mouse (rm) PDGF R β , or rmFit-3 Ligand is observed.
Source	Monoclonal Rat IgG _{2B} Clone # 460615
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse M-CSF R/CD115 Ala20-Ser511 Accession # P09581
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	RAW 264.7 mouse monocyte/macrophage cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

M-CSF receptor, the product of the *c-fms* proto-oncogene, is a member of the type III subfamily of receptor tyrosine kinases that also includes receptors for SCF and PDGF. These receptors each contain five immunoglobulin-like domains in their extracellular domain (ECD) and a split kinase domain in their intracellular region (1-4). M-CSF receptor is expressed primarily on cells of the monocyte/macrophage lineage, dendritic cells, stem cells and in the developing placenta (1). Mouse M-CSF receptor cDNA encodes a 977 amino acid (aa) type I membrane protein with a 19 aa signal peptide, a 492 aa extracellular region containing the ligand-binding domain, a 25 aa transmembrane domain and a 441 aa cytoplasmic domain. The mouse M-CSF R ECD shares >99% aa identity with rat and 60-63% aa identity with corresponding sequences in human, canine, feline and bovine M-CSF R. Activators of protein kinase C induce TACE/ADAM17 cleavage of the M-CSF receptor, releasing the functional ligand-binding extracellular domain (5). M-CSF binding induces receptor homodimerization, resulting in transphosphorylation of specific cytoplasmic tyrosine residues and signal transduction (6). The intracellular domain of activated M-CSF R binds more than 150 proteins that affect cell proliferation, survival, differentiation and cytoskeletal reorganization. Among these, PI3Kinase, P42/44 ERK and c-Cbl are key transducers of M-CSF R signals (3, 4). M-CSF R engagement is continuously required for macrophage survival and regulates lineage decisions and maturation of monocytes, macrophages, osteoclasts and DC (3, 4). M-CSF R and integrin $\alpha\beta_3$ share signaling pathways during osteoclastogenesis, and deletion of either causes osteopetrosis (7, 8). In the brain, microglia expressing increased M-CSF R are concentrated with Alzheimers $\alpha\beta$ peptide, but their role in pathogenesis is unclear (9, 10).

References:

1. deParseval, N. *et al.* (1993) *Nucleic Acids Res.* **21**:750.
2. Rothwell, V.M. and L.R. Rohrschneider (1987) *Oncogene Res.* **1**:311.
3. Chitu, V. and E.R. Stanley (2006) *Curr. Opin. Immunol.* **18**:39.
4. Ross, F.P. and S.L. Teitelbaum (2005) *Immunol. Rev.* **208**:88.
5. Rovida, E. *et al.* (2001) *J. Immunol.* **166**:1583.
6. Yeung, Y. *et al.* (1998) *J. Biol. Chem.* **273**:17128.
7. Dai, X. *et al.* (2002) *Blood* **99**:111.
8. Faccio, R. *et al.* (2003) *J. Clin. Invest.* **111**:749.
9. Li, M. *et al.* (2004) *J. Neurochem.* **91**:623.
10. Mitrasinovic, O.M. *et al.* (2005) *J. Neurosci.* **25**:4442.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.