

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse SLAM/CD150.
Source	Monoclonal Rat IgG _{2A} Clone # 459911
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse SLAM/CD150 Thr25-Pro242 Accession # Q9QUM4
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	PMA and Ca ²⁺ ionomycin-treated mouse splenocytes

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

The type I transmembrane glycoprotein Signaling Lymphocytic Activation Molecule (SLAM), also known as CD150, is the prototypic member of the SLAM subgroup of the CD2 protein family. CD2 family proteins function as adhesion molecules and modulators of the immune response (1). Mouse SLAM consists of a 218 amino acid (aa) extracellular domain (ECD) with two Ig-like domains, a 23 aa transmembrane segment, and a 78 aa cytoplasmic domain with three immunoreceptor tyrosine switch motifs (ITSM) (2). Alternate splicing generates an isoform with a substituted cytoplasmic domain (2). Within the ECD, mouse SLAM shares 58% and 83% aa sequence identity with human and rat SLAM, respectively. It is expressed as a 75 kDa molecule of which approximately 30 kDa is N-linked carbohydrate (2). SLAM is expressed on T cells, B cells, thymocytes, macrophages, dendritic cells, platelets, and hematopoietic stem cells (2-7). It is upregulated on activated B cells and CD4⁺ and CD8⁺ T cells, although it is downregulated on Th2 polarized cells (2, 3, 8). SLAM interacts homophilically with low affinity, and this interaction induces a Th0/Th1 response characterized by clonal expansion, production of IFN-γ, and increased cytolytic activity of CD8⁺ T cells (2, 3, 9-11). SLAM ligation also promotes B cell activation, allergen-induced eosinophil and mast cell activation, and macrophage responsiveness to LPS (4, 8, 12). In humans, SLAM functions as a cellular entry receptor for measles virus (13, 14).

References:

1. Ma, C.S. *et al.* (2007) *Annu. Rev. Immunol.* **25**:337.
2. Castro, A.G. *et al.* (1999) *J. Immunol.* **163**:5860.
3. Cocks, B.G. *et al.* (1995) *Nature* **376**:260.
4. Wang, N. *et al.* (2004) *J. Exp. Med.* **199**:1255.
5. Hahm, B. *et al.* (2004) *Virology* **323**:292.
6. Nanda, N. *et al.* (2005) *Blood* **106**:3028.
7. Kiel, M.J. *et al.* (2005) *Cell* **121**:1109.
8. Punnonen, J. *et al.* (1997) *J. Exp. Med.* **185**:993.
9. Mavaddat, N. *et al.* (2000) *J. Biol. Chem.* **275**:28100.
10. Aversa, G. *et al.* (1997) *J. Immunol.* **158**:4036.
11. Mehrle, S. *et al.* (2008) *Mol. Immunol.* **45**:796.
12. Wang, N. *et al.* (2006) *Am. J. Respir. Cell Mol. Biol.* **35**:206.
13. Tatsuo, H. *et al.* (2000) *Nature* **406**:893.
14. Hsu, E.C. *et al.* (2001) *Virology* **279**:9.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.