

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse IL-17 R in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human IL-17 R, recombinant mouse (rm) IL-17 RB, rmlIL-17 RD, or rmlIL-17 RC is observed. In Western blots, less than 10% cross-reactivity with recombinant human (rh) IL-17 R or rhIL-17 RB is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 657603
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse IL-17 R Ser32-Trp322 Accession # Q60943
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

IL-17 R, also known as IL-17 RA, is a widely expressed 120 kDa glycosylated type I transmembrane protein that plays a central role in inflammatory responses. IL-17 R associates with IL-17 RC to form a signaling receptor complex for IL-17 and IL-17F. IL-17 R ligation promotes T cell activation and the production of IL-6, G-CSF, SCF, and multiple proinflammatory chemokines. Within the ECD, mouse IL-17 R shares 84% and 72% aa sequence identity with rat and human IL17 R, respectively.